

**AMENDMENTS TO THE CLAIMS**

1. (Original) A process for dewatering hydrocyanic acid by distillation, which comprises distilling crude hydrocyanic acid containing from 50 to 99.9% by weight of HCN, from 0.1 to 40% by weight of water, from 0 to 15% by weight of carbon oxides and optionally from 0.01 to 1% by weight of an involatile stabilizer, at a pressure of from 1 bar to 2.5 bar, a bottom temperature of from 100°C to 130°C and a top temperature of from 25°C to 54°C, in the absence of a volatile stabilizer, in a distillation column to obtain a top draw stream containing purified, anhydrous hydrocyanic acid and carbon oxides and a bottom draw stream comprising water and, where appropriate, the involatile stabilizer.
2. (Original) A process according to claim 1, wherein the crude hydrocyanic acid comprises, as an involatile stabilizer, phosphoric acid or sulfuric acid.
3. (Original) A process according to claim 1, wherein the crude hydrocyanic acid contains no stabilizer.
4. (Currently amended) A process according to ~~any of claims 1 to 3~~ claim 1, wherein the obtained purified, anhydrous hydrocyanic acid has a water content of < 100 ppm.
5. (Currently amended) A process according to ~~any of claims 1 to 4~~ claim 1, wherein the distillation column is a bubble-cap tray column or column having structured packing.
6. (Currently amended) A process according to claim 1 ~~[to 5]~~, wherein the crude hydrocyanic acid is obtained by thermally cleaving formamide.
7. (Currently amended) A process according to ~~any of claims 1 to 6~~ claim 1, wherein the carbon oxides present in the purified, anhydrous hydrocyanic acid are stripped out using an inert gas in a downstream purification column.
8. (Currently amended) A process according to ~~any of claims 1 to 7~~ claim 1, wherein the anhydrous hydrocyanic acid is stored at from 5 to 25°C in the absence of a stabilizer over a period of from 2 to 10 days.

9. (Currently amended) A process for hydrocyanating olefins or dienes by
- a) in a first step, dewatering crude hydrocyanic acid by a process according to ~~any of claims 1 to 8~~ claim 1 and optionally storing it in the absence of a stabilizer,
  - b) reacting the anhydrous hydrocyanic acid in the absence of a stabilizer with the olefin or diene in the presence of a hydrocyanation catalyst.